


A LARGE UNTREATED CTO POPULATION

CTOs are common, but only a small proportion are treated with PCI



SUBSTANTIAL TREATMENT GAP

CTOs are present in ~15–20% of coronary angiographies, yet only ~3–6% of PCI procedures are performed for CTO.



THE MAJORITY OF CTO PATIENTS REMAIN UNTREATED OR ARE NOT CONSIDERED FOR PCI.



Improved success and safety of CTO crossing could impact a large number of patients worldwide.



REFERENCES

1. Fefer P, Knudtson ML, Cheema AN, et al. Current perspectives on coronary chronic total occlusions: the Canadian Multicenter Chronic Total Occlusions Registry. *J Am Coll Cardiol.* 2012;59(11):991–997. PMID: 22402070.
2. Othman H, Sakr K, Patel VG, et al. Percutaneous coronary intervention for chronic total occlusion—The Michigan experience: Insights from the BMC2 registry. *JACC Cardiovasc Interv.* 2020;13(11):1357–1368. PMID: 32417095.
3. van Veelen A, Elias J, Baan J Jr, et al. Incidence and outcomes of chronic total occlusion percutaneous coronary intervention in the Netherlands: data from a nationwide registry. *Neth Heart J.* 2021;29(1):4–13. PMID: 33263890.